



Metagenomics and stable isotope probing reveal the complementary contribution of fungal and bacterial communities in the recycling of dead biomass in forest soil

Rubén López-Mondéjar^{a,*}, Vojtěch Tláškal^a, Tomáš Větrovský^a, Martina Štursová^a, Rodolfo Toscan^b, Ulisses Nunes da Rocha^b, Petr Baldrian^a

^a Laboratory of Environmental Microbiology, Institute of Microbiology of the Czech Academy of Sciences, Průmyslová 595, 252 50, Vestec, Czech Republic

^b Department of Environmental Microbiology, Helmholtz Center for Environmental Research – UFZ, Permoserstr. 15, 04318, Leipzig, Germany

ARTICLE INFO

Keywords:

Decomposition
Forest soil
Microbial community
CAZyme
SIP-Metagenomic
Dead biomass

ABSTRACT

Forest soils represent important terrestrial carbon (C) pools, where C is primarily fixed in plant biomass and then is incorporated in the biomass of fungi and bacteria. Although classical concepts assume that fungi are the main decomposers of the recalcitrant organic matter within plant and microbial biomass, whereas bacteria are considered to mostly utilize simpler compounds, recent studies have shown that fungi and bacteria overlap in substrate utilization. Here, we studied the microbial contribution to the recycling of dead biomass by analyzing the bacterial and fungal communities in soil microcosms supplemented with ¹³C-labeled biomass of plant, fungal, and bacterial origin using a combination of DNA-stable isotope probing and metagenomics. Both fungi and bacteria contributed actively to the degradation of complex components of plant and microbial biomass. Specific families of carbohydrate-active enzymes (CAZyme) were involved in the degradation of each biomass type. Moreover, the analysis of five bacterial metagenome-assembled genomes indicated the key role of some bacterial genera in the degradation of plant biomass (*Cytophaga* and *Asticcacaulis*) and microbial biomass (*Herminiimonas*). The enzymatic systems utilized by bacteria are highly complex and complementary but also highly diverse among taxa. The results confirm the importance of bacteria, in addition to fungi, as decomposers of complex organic matter in forest soils.

1. Introduction

Forests represent some of the most important carbon (C) pools and sinks on Earth. Since nearly half of the C stored in these ecosystems is contained in soils, understanding the processes involved in C cycling in forest soils is essential in the current context of global climate change (Pan et al., 2011). Microorganisms are the main players involved in the recycling and turnover of soil organic matter. As such, they contribute largely to the C flow in this habitat and have the potential to influence the feedback between climate and the global C cycle (Schimel and Schaeffer, 2012). Therefore, predicting how forests will respond to future environmental conditions is impossible without understanding the roles of soil microbes in C cycling (Graham et al., 2016; Trivedi et al., 2013).

The major sources of forest soil C are comprised of the C allocated by tree roots into soil and of the C contained in the dead plant biomass in

the forms of litter and dead wood. This dead plant biomass is composed mostly of cellulose, hemicelluloses and lignin, forming a complex and recalcitrant matrix (Bomble et al., 2017). Microbial biomass represents another important pool of organic matter whose fate in the soil is far less understood. Forest soils are rich in ectomycorrhizal (ECM) and saprotrophic fungi and the decomposition of dead mycelia represents an important process for the cycling of C and other nutrients in these ecosystems (Ekblad et al., 2013; Fernandez and Koide, 2014). Dead fungal biomass is composed mainly of polysaccharides that can make up 80–90% of the total cell wall, but it also contains lipids and mannoproteins (Baldrian et al., 2013b; Fesel and Zuccaro, 2016; Free, 2013). The main components of the polysaccharide fraction include chitin, a polymer of N-acetylglucosamine units, different types of beta- and alpha-glucans, glucomannans and galactomannans. Dead bacterial biomass is considered to be equally abundant in forest soils, showing higher turnover rates than fungal biomass (Gunina et al., 2017). The

* Corresponding author.

E-mail addresses: rubenlopezmondejar@gmail.com, mondejar@biomed.cas.cz (R. López-Mondéjar).

composition of cell walls is highly diverse in bacteria (Silhavy et al., 2010). Peptidoglycan (PG), a polymer of *N*-acetylglucosamine and *N*-acetylmuramic acid units connected to chains of amino acids, is a major and universal component of bacterial cell walls (Egan et al., 2017; Scheffers and Pinho, 2005). In gram-positive bacteria, PG is densely functionalized with other polymers. Cell-wall glycopolymers such as teichoic, teichuronic and teichulosonic acids, which are attached either to the PG or to membrane lipids, are the most abundant (Brown et al., 2013; Schaffer and Messner, 2005; Weidenmaier and Peschel, 2008). Apart from PG and cell-wall glycopolymers, the bacterial cell wall includes proteins, glycosyl 1-phosphates and other sugar-containing polymers such as arabinogalactan, lipomannan and lipoarabinomannan (Hamed and Poorinmohammad, 2017). Furthermore, gram-negative bacteria contain lipopolysaccharides and lipoproteins in their outer membranes (Silhavy et al., 2010). Many bacteria also produce a range of chemically different extracellular polysaccharides, which can be utilized as C sources by other microorganisms in soils (Bazaka et al., 2011; Mishra and Jha, 2013; Wang et al., 2015).

The turnover of carbohydrates in plant and microbial biomass can be tracked by analyzing the microbial enzymes that take part in the C turnover—the carbohydrate-active enzymes (CAZymes) (Žifčáková et al., 2017). CAZymes, classified into a hierarchy of families based on their structure and function, act on oligosaccharides, polysaccharides and glycoconjugates (Lombard et al., 2014). Among the CAZymes, glycoside hydrolases (GHs), which hydrolytically cleave the glycosidic bonds within carbohydrates or between a carbohydrate and a non-carbohydrate moiety, are the most important in decomposition. In this sense, cellulases, β -glucosidases and hemicellulases such as endoxylanases, β -xylosidases, xyloglucanases, endomannanases, mannosidases, fucosidases, and arabinosidases from several GH families are the main enzymes that degrade plant biomass (Bomble et al., 2017). Beside them, lytic polysaccharide monooxygenases (LPMOs), classified as enzymes with auxiliary activities (AA) in the CAZy database, have also been found to play an important role in the degradation of cellulose (Vaaje-Kolstad et al., 2017). In addition, several AA families including peroxidase, oxidoreductase and laccase activities participate either directly or indirectly in the degradation of lignin (Levasseur et al., 2013). Finally, carbohydrate esterases (CEs) from several families participate in the decomposition of hemicelluloses. In the case of fungal biomass, chitinases and *N*-acetylglucosaminidases from three GH families are involved in the degradation of chitin, and glucanases from several GH families, which degrade glucans, are highlighted as main players involved in its degradation. The lysozymes and PG lytic transglycosylases are important enzymes involved in the degradation of PG in bacterial biomass. Catalytically active CAZymes may contain carbohydrate-binding modules (CBMs), which are essential for effective hydrolysis because they mediate binding to cellulose, xylan, chitin or other carbohydrates (Donohoe and Resch, 2015). Many GH families include enzymes that are structurally similar but have wider substrate specificity, and associating one family to the degradation of one type of compound is not always easy (Nguyen et al., 2018). Moreover, the complex, diverse and not fully characterized composition of dead biomass in forest soils, especially in the case of fungal and bacterial biomass, may entail the implication of more CAZyme families than those currently proposed.

For a long time, fungi were assumed to be the major decomposers of complex organic matter in forest soils due to their filamentous nature, which allows them to colonize substrates efficiently, their ability to produce a rich battery of extracellular enzymes and their limited requirements of N, which is rather rare in cell wall biopolymers. This assumption led to underestimation of the role of bacteria in decomposition, and bacteria were typically expected to target simple substrates (de Boer et al., 2005; Rousk and Frey, 2015). Different studies have indicated that bacteria play a more important role in the transformation and mineralization of organic matter and contribute significantly to decomposition in forest soils (Eichorst and Kuske, 2012; Stursová et al.,

2012; Verastegui et al., 2014). The high percentage of bacteria that potentially decompose cellulose found in forest soil and the high frequency of genes involved in the degradation of structural plant polysaccharides found in bacterial genomes support that the involvement of bacteria in plant biomass decomposition is relatively common (Berlemont and Martiny, 2015; López-Mondéjar et al., 2016a; Wilhelm et al., 2019). In addition, analyses of forest soil metatranscriptomes show significant contribution of bacteria to CAZyme production (Hesse et al., 2015; Lladó et al., 2019; Žifčáková et al., 2017). Moreover, Brabcova et al. (2016) showed that decomposing mycelium in forest soil presents hotspots of bacterial abundance, maintaining bacterial over fungal decomposers. In our previous work, we demonstrated that both fungi and bacteria are involved in the assimilation and mineralization of C from complex sources existing in soil. In addition, we showed that fungi may be better suited for the utilization of plant biomass, whereas most bacteria prefer microbial biomass (López-Mondéjar et al., 2018).

The aim of this study was to describe the enzymatic toolbox used for the decomposition of various biomass types by forest soil bacteria and fungi. To accomplish this, we prepared soil microcosms with the addition of ^{13}C -labeled biomass of plant, fungal, and bacterial origin. We used DNA-SIP and metagenomics to analyze the enzymatic tools of microbial decomposers. We hypothesized that although the CAZyme pool will be different for each type of biomass, fungi and bacteria will encode similar CAZyme families involved in the degradation of biomass of the same origin. Additionally, in line with the preference of fungi for plant biomass, we hypothesized that the number of fungal CAZymes involved in the degradation of plant biomass will be higher than the number of those targeting microbial biomass. Importantly, this study also provides a comprehensive answer about the CAZyme families involved in the degradation of various biomass types, including the involvement of minor CAZy families that might have been overlooked so far.

2. Material and methods

2.1. Sample collection

Soil was collected from the organic horizon of a sessile oak (*Quercus petraea*) forest in the Xaverovský Háj Natural Reserve in the Czech Republic. Previously, this site has been studied with respect to the composition of microbial communities and their seasonal changes and the activity of extracellular enzymes related to the decomposition process (Šnajdr et al., 2008; Baldrian et al., 2010, 2013a; Voříšková et al., 2014; López-Mondéjar et al., 2015). This study used the samples collected previously, which were incubated as described in the study of López-Mondéjar et al. (2018). Briefly, soil was sieved and preincubated at 10 °C for 48 h and allocated in 100-ml flasks containing 5 g of soil and 0.08 g of different ^{13}C -labeled substrates: ^{13}C -glucose (99 atom% ^{13}C), ^{13}C -cellulose (from *Zea mays*, 97 atom% ^{13}C), ^{13}C -hemicellulose (from *Zea mays*, 97 atom% ^{13}C), ^{13}C -plant biomass (from ground maize leaves, 97 atom% ^{13}C), ^{13}C -bacterial biomass (from *Streptomyces* sp. PR6, prepared by cultivation in media with ^{13}C -glucose as sole C source) and ^{13}C -fungal biomass (from *Phanerochaete velutina* PV29, prepared by cultivation in media with ^{13}C -glucose as sole C source). Microcosms were slightly moistened with water to reach 60% water content and incubated at 10 °C in the dark for 3 weeks (21 days). After this, three microcosms per treatment were harvested and the material was frozen at -80 °C, freeze-dried and stored at -40 °C.

2.2. DNA extraction and sequencing

DNA was extracted with the FastDNA Spin Kit for Soil (MP Biomedicals) in triplicate and purified with the GeneClean Turbo Kit. Three micrograms of total DNA was used for isopycnic centrifugation in a cesium trifluoroacetate (CsTFA) solution to separate the labeled and unlabeled fractions, as shown in López-Mondéjar et al. (2018). After centrifugation, the labeled fractions representing the ^{13}C -DNA were

pooled for each microcosm. Due to the small amount of DNA recovered after centrifugation, the labeled DNA from triplicates from the same substrate was also pooled, and the total DNA was used for metagenome sequencing. DNA libraries were prepared using the KAPA Hyper Prep Kit (Roche) according to the manufacturer's instructions. Metagenome libraries were sequenced on an Illumina HiSeq 2000 to generate 250-base paired-end reads. In total, eight metagenomes were sequenced, including the ^{13}C -labeled DNA isolated after 21 days of incubation of soil with ^{13}C -substrate (glucose (GL), cellulose (CE), hemicellulose (HE), plant biomass (PB), bacterial biomass (BB) and fungal biomass (FB)) and the DNA from the control microcosms (with no substrate addition) at both 0 (C1) and after 21 (C2) days of incubation.

2.3. Metagenome assembly

Reads from all the metagenome libraries were processed together in the same way as originally described in Žifčáková et al. (2016). Briefly, the reads were quality trimmed by removing adapters, filtered by base call quality, and normalized. Errors were trimmed by removing low abundance fragments of high coverage reads. The paired-end assembly of the remaining reads was performed with the Velvet assembler (v 1.2.10) (Zerbino and Birney, 2008) using odd k-mer lengths ranging from 33 to 63. Resulting assembled contigs were merged using CD-HIT v4.6 (Li and Godzik, 2006; Fu et al., 2012) and minimus2 Amos v3.1.0 (Sommer et al., 2007). Sequence data of all contig sequences (whole metagenome) were deposited in the MG RAST database under the dataset number mgs446518.

Annotation of contigs was performed using both MG RAST (Meyer et al., 2008) and an in-house fungal-predicted protein database (FPPD), as described by Žifčáková et al. (2017). The taxonomic classification for each contig was retrieved from that of the two databases which showed lower bitscore values of the best hit.

2.4. CAZyme annotation

The annotation of CAZymes in the metagenome contigs was carried out after gene calling using the pipeline dbCAN (Yin et al., 2012), followed by manual curation considering the alignments to the sequences in the CAZy database (February 2018) (Table 1). To assess the abundance in the metagenome, individual sequence reads from each sample were mapped onto contigs identified as CAZyme using bowtie 2.2.1 (Langmead et al., 2009), with the default settings of end to end alignment—sensitive. Data were expressed as: per base coverage = read count \times read length/contig length to calculate gene abundance, as previously described (Žifčáková et al., 2017). R software (RCoreTeam, 2019) was used for statistical analysis. Differences in gene abundance in each metagenome and the controls were tested using the exact Fisher test (Gharechahi and Salekdeh, 2018). Differences at $P < 0.05$ were considered statistically significant.

2.5. Recovery and analysis of metagenome assembled genomes (MAGs)

The recovery of metagenome-assembled genomes (MAGs) was performed as follows. First, quality-controlled reads (see earlier) were assembled into scaffolds using IDBA-UD (Peng et al., 2012). Second, the scaffolds were assigned as eukaryotic or prokaryotic using the EUKREP pipeline (West et al., 2018). No significant number of scaffolds was assigned as eukaryotic; hence, we did not attempt to recover the eukaryotic bins. Prokaryotic scaffolds were binned using ABAWACA (<https://github.com/CK7/abawaca>), Maxbin2 (Wu et al., 2015), MetaBAT2 (Kang et al., 2019), and CONCOCT (Alneberg et al., 2014) and further refined with the DAS Tool refinement method (Sieber et al., 2018). Completeness and contamination values for each bin were determined using CheckM (Parks et al., 2015). Bins were considered MAGs when the quality score was above 50, as defined by Parks and collaborators (Parks et al., 2017). Briefly, this quality score is defined by the percentage of

Table 1

List of the main CAZyme families encoding the enzymatic activities involved in the degradation of several compounds presented in plant and microbial biomass according to CAZy (<http://www.CAZy.org>).

Origin	Compound	CAZyme families
Plant biomass	cellulose	GH1 (β -glucosidase), GH3 (β -glucosidase), GH5 (β -glucosidase/endoglucanase), GH6 (cellobiohydrolase), GH7 (reducing end-acting cellobiohydrolase), GH8 (endoglucanase/endoxylanase), GH9 (endoglucanase), GH12 (endoglucanase), GH45 (endoglucanase), GH48 (reducing end-acting cellobiohydrolase/endoglucanase), GH116 (β -glucosidase), AA9 (lytic polysaccharide monooxygenase), and AA10 (lytic polysaccharide monooxygenase)
	hemicellulose	GH2 (β -galactosidase/ β -glucuronidase), GH10 (endoxylanase), GH11 (endoxylanase), GH16 (xyloglucanase/endoglucanase), GH26 (endomannanase), GH30 (endoxylanase/ β -1,6-glucanase/ β -xylosidase), GH36 (α -galactosidase), GH39 (β -xylosidase/ α -L-arabinofuranosidase), GH43 (β -xylosidase/endoxylanase), GH44 (xyloglucanase/endoglucanase), GH51 (α -L-arabinofuranosidase), GH52 (β -xylosidase), GH54 (α -L-arabinofuranosidase), GH62 (α -L-arabinofuranosidase) GH67 (xylan α -1,2-glucuronidase), GH74 (xyloglucanase), GH95 (α -L-fucosidase/ α -L-galactosidase), GH115 (xylan α -1,2-glucuronidase), GH120 (β -xylosidase), GH131 (exo- β -1,3/1,6-glucanase/endo- β -1,4-glucanase) and CE1, CE2, CE3, CE4, CE5, CE6, CE7, CE12, CE15, CE16 (acetyl xylan esterases)
	lignin	AA1 (laccase), AA2 (peroxidase), AA3 (oxidase), AA4 (oxidase), AA5(oxidase), AA6 (1,4-benzoquinone reductase),
Fungal biomass	chitin	GH18 (chitinase), GH19 (chitinase), GH20 (N-acetyl β -glucosaminidase) and AA11 (lytic polysaccharide monooxygenase)
	glucans	GH17 (endo-1,3- β -glucanase), GH55 (exo- β -1,3-glucanase/endo-1,3- β -glucanase), GH64 (endo-1,3- β -glucanase), GH81 (endo-1,3- β -glucanase), and GH128 (endo-1,3- β -glucanase)
Bacterial biomass	peptidoglycan	GH22 (lysozyme), GH23 (lysozyme/peptidoglycan lytic transglycosylase), GH24 (lysozyme), GH25 (lysozyme), GH73 (peptidoglycan hydrolase with endo- β -N-acetylglucosaminidase specificity), GH102 (peptidoglycan lytic transglycosylase), GH103 (peptidoglycan lytic transglycosylase), GH104 (peptidoglycan lytic transglycosylase) and GH108 (lysozyme)

completeness of a bin minus five times the percent contamination. The taxonomy was assigned to MAGs using GTDB version v0.2.2 (Parks et al., 2018) and the Microbial Genomes Atlas (MiGA) webserver (Rodríguez et al., 2018). MAGs were compared with all available genomes of bacteria belonging to the same family using *anvi'o* v5.5 (Eren et al., 2015) and the average nucleotide identity (ANI) was calculated. Average amino acid identity (AAI) was computed using the CompareM (v0.0.23) AAI workflow (D. H. Parks, unpublished materials, <https://github.com/dparks1134/CompareM>). The CAZyme content of these genomes was predicted using dbCAN2 (Zhang et al., 2018) followed by manual curation using the CAZy database.

3. Results

3.1. Total diversity of the CAZyme pool in the metagenome

In total, 132,197 CAZymes were identified from the 7.2 million predicted proteins of the whole metagenome (1.84% of genes), of which 26.0% and 42.6% were assigned to bacteria and fungi, respectively, and the rest (31.4%) were unassigned. The bacterial CAZymes belonged to 233 families, including 95 GHs, 4 AAs, 14 CEs, 17 Polysaccharide Lyases

(PLs), 49 CBMs and 54 GlycosylTransferases (GTs), and the fungal CAZymes were assigned to 209 families, including 84 GHs, 10 AAs, 11 CEs, 9 PLs, 32 CBMs and 63 GTs (Figs. S1, S2, S3 and S4). Bacterial GHs accounted for 9.1% of all CAZymes with the highest diversity in the families GH13 (amylase/ α -glucosidase/trehalase), GH109 (α -N-acetylgalactosaminidase), GH3 (β -glucosidase), GH23 (lysozyme/peptidoglycan lytic transglycosylase) and GH15 (glucoamylase/glucodextranase). Fungal GHs accounted for 16.2% of all CAZymes. Similar to bacteria, GH3, GH13 and GH15 were the most diverse families, in addition to GH43 (β -xylosidase/endoxylanase) and GH31 (α -glucosidase/ α -xylosidase). The number of fungal genes encoding AAs and CEs was twice as high (1.7 and 8.5%, respectively) as bacterial genes (0.5 and 4.5%), but similar for PLs (0.3 vs 0.4%), GTs (7.7 vs 8.1%) and CBMs (2.7 vs 2.7%).

In general, most of the fungal CAZymes were assigned to Ascomycota (55.9%), Basidiomycota (21.2%) and Mucoromycota (9.3%). Among them, Ascomycota showed a higher number of contigs encoding GHs, AAs, CBMs, CEs, PLs and GTs (Fig. 1). In the case of bacteria, the contigs encoding CAZymes were mainly assigned to Proteobacteria (28.4%), Actinobacteria (25.7%), Acidobacteria (20.3%) and Bacteroidetes (16.4%). These four phyla also showed the highest numbers of CAZymes of all six classes (Fig. 1).

3.2. CAZyme pools in the fungal and bacterial communities

Relative abundances of genes assigned to CAZyme families in the ^{13}C -enriched metagenomic DNA from microcosms containing ^{13}C -supplemented substrates were compared with the abundance in the

metagenome of the control samples individually. In total, the percentage of CAZyme families that were significantly increased in at least one of the ^{13}C -supplemented treatments was higher in fungi (69%) than in bacteria (49%) (Fig. 2, Fig. 3). The increase in the relative abundance of certain CAZy families may indicate a relative increase in the abundance of microorganisms utilizing carbon from certain sources. When ^{13}C -cellulose was added to the soil, more CAZyme families were enriched in the fungal community than in the bacterial community (24 vs 18, respectively). On the other hand, the bacterial community showed more CAZyme families increasing upon the addition of ^{13}C -fungal biomass (52 vs 39) and ^{13}C -bacterial biomass (52 vs 37). For ^{13}C -plant biomass and ^{13}C -hemicellulose, the numbers were similar for bacteria and fungi; 52 and 53 in plant biomass, 67 and 69 in hemicellulose, respectively.

Both fungal and bacterial communities were enriched in numerous CAZymes known to be involved in the degradation of plant, fungal and bacterial biomass (Figs. 2 and 3). The CAZyme pool in the plant and microbial dead biomass metagenomes was composed of numerous well-known fungal- and bacterial-encoded CAZyme families including cellulases, glucosidases, xylanases, xylosidases, mannosidases, galactosidases, glucanases, xyloglucanases, glucuronidases, chitinases, hexosaminidases, lysozymes, acetyl xylan esterases and CBMs binding several polysaccharides.

3.3. CAZyme families involved in the degradation of plant, fungal and bacterial biomass

Although many of the CAZyme families were significantly increased in all of the ^{13}C metagenomes (Figs. 2 and 3), several of them were only




Fig. 1. Diversity of genes in the whole metagenome encoding CAZymes assigned to bacterial (A) and fungal (B) phyla. GH: Glycoside Hydrolases, AA: Auxiliary Activities, CBM: Carbohydrate-Binding Modules, CE: Carbohydrate Esterases, PL: Polysaccharide Lyases, GT: GlycosylTransferases.






Fig. 2. Relative abundance of genes assigned to CAZyme families of fungi in the ¹³C-incorporating microbial community compared to the control. The values indicate fold enrichment in the ¹³C-DNA of each treatment/total DNA of control without substrate addition. Genes significantly more abundant in treatment than in control are shown in yellow-red, genes significantly less abundant are shown in light and dark blue, white squares indicate no significant difference between control and treatment. Abbreviations indicate ¹³C substrate addition of FB: fungal biomass; BB: bacterial biomass; PB: plant biomass; CE: cellulose; and HE: hemicellulose. Only CAZy families with relative abundance significantly different from control in at least one of the substrates are shown. Families containing enzymes degrading cellulose or hemicellulose have a green mark, fungal biomass blue and bacterial biomass purple (source: www.cazy.org). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

enriched in a specific type of dead biomass (Fig. 4, Fig. 5). In general, the fungal community contained a richer set of CAZyme families that were specifically increased upon the addition of ¹³C-labeled plant biomass and its components. The bacterial community also contained several CAZyme families that increased after the addition of plant biomass, but this community was particularly rich in CAZymes that increased after the addition of ¹³C-labeled bacterial and fungal biomass.

Regarding the fungal CAZymes involved in the degradation of plant biomass, we found several families encoding enzymes that degrade cellulose and hemicelluloses (Fig. 4). In this sense, CAZyme families containing β-glucosidases (GH1 and GH3), endoglucanases (GH5, GH9, and GH45), cellobiohydrolases (GH7), endoxylanases (GH10 and GH11), other hemicellulases (GH62, GH67 and GH131), LPMOs and oxidases (AA1, AA7 and AA9), and CBM families binding to plant



(caption on next page)

Fig. 3. Relative abundance of genes assigned to CAZyme families of bacteria in the ¹³C-incorporating microbial community compared to the control. The values indicate fold enrichment in the ¹³C-DNA of each treatment/total DNA of control without substrate addition. Genes significantly more abundant in treatment than in control are shown in yellow-red, genes significantly less abundant are shown in light and dark blue, white squares indicate no significant difference between control and treatment. Abbreviations indicate ¹³C substrate addition of FB: fungal biomass; BB: bacterial biomass; PB: plant biomass; CE: cellulose; and HE: hemicellulose. Only CAZY families with relative abundance significantly different from control in at least one of the substrates are shown. Families containing enzymes degrading cellulose or hemicellulose have a green mark, fungal biomass blue and bacterial biomass purple (source: www.cazy.org). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)




Fig. 4. Fungal CAZyme families enriched after addition of each one of the different ¹³C substrates: in cellulose, in plant biomass or hemicellulose, in bacterial biomass, in fungal biomass and in both types of microbial biomass and their known catalytic properties. Abundance represents the total number of CAZymes of that family found in the whole metagenome. In bold, families showing activities directly involved in the degradation of that substrate.




Fig. 5. Bacterial CAZyme families enriched after addition of each one of the different ¹³C substrates: in cellulose, in plant biomass or hemicellulose, in bacterial biomass, in fungal biomass and in both types of microbial biomass and their known catalytic properties. Abundance represents the total number of CAZymes of that family found in the whole metagenome. In bold, families showing activities directly involved in the degradation of that substrate.

biomass components (CBM1, CBM6, CBM20, CBM21) increased in abundance when plant biomass, cellulose or hemicellulose were added to soil. Interestingly, we found other families that were not related to the

degradation of plant biomass and their components, but encoding chitinases, glucanases, chitosanases or glucanosyltransferases (e.g. GH72, GH75, GH79) and binding to chitin (CBM5 and CBM18). In the bacterial community, plant biomass amended soil was enriched in different CAZY families than those of fungi (Fig. 5). Although most of the families are also known to be involved in plant polysaccharide degradation, such as those encoding endoglucanases (GH5 and GH9), β -glucosidases (GH116), endomannanases (GH26), endoxyylanases (GH30), xyloglucanases (GH44 and GH115), acetyl xylan esterases (CE2, CE4, CE7, CE8, CE14 and CE15) and CBMs binding cellulose (CBM3), the rest of the CAZymes belonged to families that have not been shown to participate in plant biomass degradation. The families GH4 (α -galactosidase/6-phospho- β -glucosidase/maltose-6-phosphate glucosidase), GH109 (α -N-acetylgalactosaminidase), AA6 (1,4-benzoquinone reductase) and the domains CBM32 (binding to D-galactose and N-acetyl-D-galactosamine) and CBM50 (binding to N-acetylglucosamine) were abundant in the metagenome.

Meanwhile, most of the fungal CAZymes enriched in ^{13}C -bacterial biomass were assigned to families with potential activity in bacterial biomass degradation. The roles of several families of bacterial CAZymes in the decomposition of bacterial biomass are still unclear (Figs. 4 and 5). Regarding the degradation of ^{13}C -fungal biomass, both bacterial and fungal CAZymes belonged to several families with uncertain roles in the process, such as the family GH32, whose abundance increased in both the fungal and bacterial communities.

3.4. Bacterial taxa and CAZymes degrading the main types of dead biomass in forest soil

Six MAGs were obtained from the metagenomes, but only 5 of them showed acceptable quality for further analyses (Table S1). Four MAGs belonged to Proteobacteria, and the other two belonged to Bacteroidetes. Since databases only classified the MAGs up to the family level, we compared MAGs with the available genomes in the respective family. The results showed that MAG_2 and MAG_6 were most similar to *Asticcacaulis benevestitus* DSM 16100 (76.3% and 75.9% ANI, respectively), *Asticcacaulis biprosthecium* C19 (75.11% and 75.0% ANI, respectively) and *Asticcacaulis excentricus* CB 48 (74.58% and 74.15% ANI, respectively) (Fig. S5). MAG_3 showed the highest similarity to *Cytophaga aurantiaca* DSM 3654 (77.25% ANI) and to *Cytophaga hutchinsonii* ATCC 33406 (75.67% ANI) (Fig. S6). MAG_4 and MAG_5 showed the highest similarity to *Herminiimonas arsenicoxydans* DSM 17148 (73.35% and 73.52% ANI, respectively) and to *Collimonas fungivorans* NCCB 100033 (73.06% and 73.07% ANI, respectively) (Fig. S7). The same similarity results were found when calculating the AAI values for all the MAGs (Supplementary File S1).

MAG_2, MAG_3 and MAG_6 showed the highest abundance on ^{13}C -plant biomass, while MAG_4 and MAG_5 were most abundant in the ^{13}C -bacterial biomass metagenome (Fig. S8). These results corresponded with the composition of their CAZY gene sets (Table S2). For example, MAG_3 presented numerous genes of families involved in plant biomass degradation, such as endoglucanases (GH9: 5 genes, GH5: 4 genes), hemicelluloses (GH26: 2 genes, GH30: 2 genes, GH44: 1 gene, GH74: 2 genes) and esterases (CE2: 1 gene, CE4: 6 genes and CE14:1 gene). The CAZY composition of the other two MAGs abundant in plant biomass amended soil was different. Both MAG_2 and MAG_6 also encoded genes of the GH9 (one gene each) and GH5 (two genes each) families and CEs, although to a lesser extent than MAG_3. However, both genomes contained several genes belonging to the hemicellulolytic families, such as GH39 (1 and 2 genes), GH42 (1 and 2 genes), GH51 (3 genes), GH67 (1 gene) and GH115 (2 genes), which were not detected in MAG_3. MAG_4 and MAG_5 increased upon bacterial biomass addition, and none of the CAZY families with known involvement in bacterial biomass degradation were found, with the exception of GH108 (lysozyme) in MAG_4. On the other hand, MAG_4 and MAG_5 contained genes from the families GH36 (α -galactosidase), GH73 (peptidoglycan hydrolase with endo- β -N-

acetylglucosaminidase specificity) and GH94 (phosphorylases), which were highly enriched in the ^{13}C -bacterial biomass metagenome (approximately 2 and 5 times higher abundance than in ^{13}C -plant biomass (Fig. 3)). Interestingly, other families only found in the ^{13}C -bacterial biomass metagenome, such as GH39 (β -xylosidase/ α -L-arabinofuranosidase) and GH105 (unsaturated glucuronyl/galacturonidase), were present in the genomes of MAG_2 and MAG_6, despite these bins were not abundant in the BB metagenome.

4. Discussion

Using DNA-SIP and metagenomics, our study revealed that fungi and bacteria that utilize C from plant and microbial biomass possess numerous CAZymes involved in the degradation of both types of biomass. This finding supports the recent view that the roles of fungi and bacteria as primary consumers of complex substrates are important (Kramer et al., 2016; Rousk and Frey, 2015; Žifčáková et al., 2017).

Despite the possibility that fungal CAZY predictions in metagenomes are underestimated due to introns in sequences, which make gene calling less reliable (Fierer et al., 2012; Pold et al., 2016), we still found that the number of CAZymes assigned to fungi was larger than that of bacteria. The presence and abundance of numerous well-known families encoding cellulases and hemicellulases in bacteria confirm their role in the degradation of plant-derived biomass. These results add to the list of previous evidence on the active contribution of bacteria in the decomposition of cellulose and hemicellulose (López-Mondéjar et al., 2016a; Štursová et al., 2012; Wilhelm et al., 2019) and confirm that the increase in labeled bacterial biomass after incubation is due to the actual degradation of plant biomass by hydrolytic enzymes and not just a result of mutualistic feeding by cross-feeders in SIP experiments (López-Mondéjar et al., 2018).

Contrary to our first hypothesis, the pool of CAZymes for degrading the diverse types of biomass was distinct in fungi and bacteria. In this sense, some CAZY families seem to be specific for the decomposition of plant biomass and its components in fungi and others in bacteria. While bacteria contain more esterases, fungi possess more oxidases, laccases and monooxygenases. The presence of these complex fungal enzymatic systems for degrading plant biomass, which include not only GHs but also LPMOs, may explain why fungi seem better suited to utilize plant-derived compounds (López-Mondéjar et al., 2018). Bacteria contained more CBMs than fungi, which points at the importance of substrate binding. Unlike fungi that produce mostly extracellular enzymes, cell-bound enzymes play a more important role in the degradation of organic matter in soil bacteria (Lasa et al., 2019). Therefore, the presence of CBMs in bacterial lytic enzymes allows the direct association of the bacterial cell with the target polysaccharide, improving the efficacy of the lytic enzymes and increasing the competitive exclusion of non-cellulolytic opportunists (Donohoe and Resch, 2015). Interestingly, after the addition of plant biomass and its components, fungi also showed high abundance of specific CBM families assigned to chitin- and glucan-binding (typical fungal cell wall components), which, however, were not increased in the fungal biomass metagenome. Moreover, the addition of these plant-derived substrates also increased the abundance of GHs from families potentially encoding chitinases, glucanases, chitosanases and glucanosyltransferases (such as GH18, GH71, GH72 and GH75). The abundance of fungal enzymes that target and bind to fungal cell wall components may indicate a strong competition among fungi for plant-derived substrates. These competitive fungal-fungal interactions for resources among different taxonomic groups often occur in soils (Boddy and Hiscox, 2016; Woodward and Boddy, 2008), which is especially important between different fungal guilds (Fernandez and Kennedy, 2016). The fact that these families were not increased in the fungal biomass metagenome may be explained by the differences in the attractiveness of the living and dead fungal biomass as a target for attack, such as the nutrient content or the presence of secondary compounds (Fernandez et al., 2016) and by the reportedly minor role of fungi in the

decomposition of dead fungal biomass (Brabcová et al., 2016; López-Mondéjar et al., 2018).

In accordance with our second hypothesis, we demonstrated that fungal communities encode more specific CAZymes that degrade plant biomass, while bacterial communities are richer in CAZymes that target microbial biomass. As mentioned above, when cellulose was added to soil more fungal CAZymes were increased, including not only endoglucanases as in the bacterial community but also LPMOs and CBMs for cellulose binding. Similar results were presented by Žifčáková et al. (2017), who found that fungi are the major producers of CAZymes involved in lignocellulose degradation in spruce forests, while the transcription of CAZymes involved in the degradation of bacterial and fungal cell walls was increased in bacteria. Our results are in accordance with previous studies supporting the role of fungi as the agents primarily responsible for the transformation of plant-derived carbon in terrestrial ecosystems and reinforce the evidence that bacteria are the main decomposers of mycelia in litter and soils (Bhatnagar et al., 2018; Brabcová et al., 2016; Tláškal et al., 2016; Voříšková et al., 2014). Microbial biomass represents a more readily decomposable substrate than lignocellulose, and bacteria have been shown to dominate the initial phase of dead fungal biomass decomposition (Brabcová et al., 2018). As noted previously, the different nitrogen content between microbial and plant biomass could explain the high abundance of bacteria in the decomposition of microbial biomass (López-Mondéjar et al., 2018). Bacteria contain more N in their biomass than fungi, and specialization to the decomposition of N-rich biomass may reflect their higher nutritional demand (Wallenstein et al., 2006).

Linking genes to environmental processes is vital for understanding the role of microbes in ecosystems (Graham et al., 2016). In the case of CAZyme families, it is possible to link genes with targeted substrates and thus with a specific part of the soil decomposer food web (Nguyen et al., 2018). The links between the CAZyme families, catalytic functions and substrates demonstrated here, were, however, not always entirely clear. While as much as 70% of the CAZyme families that were enriched after addition of plant-derived biomass are known to target plant biomass components, in the case of fungal or bacterial biomass, the percentage of CAZyme families known to target them was smaller. Despite some CAZyme families showing unique substrate specificity, most families have wide substrate specificity; moreover, this substrate specificity is often unknown for several families (Nguyen et al., 2018). Although gene enrichment after substrate addition may indicate the involvement of a gene product in the processing of the substrate, none of the two highly abundant MAGs on bacterial biomass contained any of the known BB-specific genes, while they contained other families such as GH73 and GH108, both encoding lysozymes. Although these two families were enriched in bacterial biomass, they were not specific and appeared on plant and fungal biomass amended soil. Previous studies using SIP showed that the degradation of some bacterial components in soil is mainly carried out by bacteria with low identities to known species, mostly from uncultured bacteria from the *Planctomycetes*, *Armatimonadetes* and the Candidate Phyla Radiation (CPR) groups (Wang et al., 2015). Most of these taxa are still poorly represented in the databases with only a low number of available sequences, which may hinder the biochemical characterization of the genes potentially involved in the degradation of microbial biomass (Lloyd et al., 2018).

After demonstrating that bacteria incorporate significant amounts of C from dead plant biomass (López-Mondéjar et al., 2018), the enrichment of bacterial CAZyme families targeting plant biomass that we showed here confirms that bacteria indeed contribute to plant biomass degradation. Two potentially different enzymatic systems for the degradation of plant biomass were observed across MAGs: the first was represented by *Cytophaga* (MAG_3) and the second was represented by *Asticcacaulis* (MAG_2 and MAG_6). Although some of the genes encoding hydrolytic enzymes were common, the genome of *Cytophaga* contained more genes potentially encoding endoglucanase, endomannanase, endoxylanase and xyloglucanase activities. *Asticcacaulis* contained fewer

endoglucanases, but more genes encoding debranching hemicellulases, such as glucuronidases, α -fucosidases, β -galactosidases, α -L-arabinofuranosidase, β -glucosidases and β -xylosidases. In line with these differences, *Asticcacaulis* was more abundant on hemicellulose. *Cytophaga* is a common cellulolytic soil bacterium that has a unique mechanism for cellulose degradation (López-Mondéjar et al., 2019). As previously found by other authors, the recovered genome did not contain any cellobiohydrolases or LPMOs, showing that the system for cellulose degradation used by this bacterium is still poorly understood (Wang et al., 2017; Zhu et al., 2016). *Asticcacaulis* spp. have been isolated from soils, and their utilization of cellulose has been demonstrated using SIP (Eichorst and Kuske, 2012; Štursová et al., 2012). It was recently demonstrated that *Asticcacaulis* assimilated ^{13}C from both cellulose and hemicellulose added to forest soil, indicating its high abundance and its ability to adhere to lignocellulose, suggesting its role in the decomposition of cellulose and hemicellulose (Wilhelm et al., 2019). Our previous results also showed that the genera *Asticcacaulis* and *Cytophaga* are specialists utilizing plant-derived compounds. Both genera were abundant in the bacterial community when ^{13}C -plant biomass or ^{13}C -cellulose was added to soil, while *Asticcacaulis* was also abundant in samples with ^{13}C -hemicellulose (López-Mondéjar et al., 2018). The presence of structurally variable enzymatic systems for decomposing cellulose and hemicellulose among different cellulolytic bacteria has been previously reported in several taxa abundant in forest soils (López-Mondéjar et al., 2016a, 2016b). Here, the complementary nature of the two enzymatic systems of these abundant and specialist bacterial taxa, one targeting the backbones and the other targeting the branches of lignocellulose, confirm that in natural environments, plant biomass is degraded by the cooperation of complex microbial communities rather than by a single species, as previously assumed (Cragg et al., 2015; Cavaliere et al., 2017; López-Mondéjar et al., 2019).

Our results also indicate the importance of *Herminiimonas* in the degradation of dead microbial biomass, which is used by this genus as a C source (López-Mondéjar et al., 2018). To date, only few species of this genus have been described, mainly those isolated from water but also from urban and contaminated soils (Koh et al., 2017; Sahin et al., 2010). Previous studies showed that *Herminiimonas* is abundant in forest soils (Bárta et al., 2017), where it accumulates C from cellulose and plant biomass (López-Mondéjar et al., 2018; Štursová et al., 2012). This ability can be explained by the presence of several genes involved in plant biomass degradation in the MAGs related to *Herminiimonas*, although in smaller numbers than in the MAGs related to *Cytophaga* and *Asticcacaulis*. For example, three potential endoglucanases from families GH9 and GH5 were found both in MAG_4 and MAG_5 but no endoxylanases or other hemicellulose backbone-degrading enzymes. However, these genomes encoded numerous genes from the families GH23, GH24, GH73 and GH108, potentially encoding lysozymes for degrading bacterial cell walls, which may explain their enrichment after bacterial biomass addition. Moreover, both MAGs contained genes encoding a β -1,2-glucanase from the family GH144, which is involved in the degradation of glucans naturally present in some bacteria (Abe et al., 2017). Recently, *Herminiimonas* has been reported to be involved in the degradation of mycelial necromass in forest soil (Sukdeo et al., 2019). However, the analysis of our two genomes revealed only one (in MAG_5) and two (in MAG_4) genes encoding for chitinases and no *N*-acetylglucosaminidases for chitin utilization; moreover, none of the CAZy families enriched on fungal biomass were present. In accordance with our previous results, *Herminiimonas* appears to be a generalist decomposer that is able to use substrates of various origins (López-Mondéjar et al., 2018).

In summary, the current study confirms that both fungi and bacteria are involved in the recycling of dead biomass in forest ecosystems and in the assimilation and mineralization of C of both plant and microbial origin. In addition to the proven role of fungi in the decomposition of plant biomass, members of the bacterial community appear to be important players in the degradation of plant-derived compounds by using structurally variable enzymatic systems. Moreover, the

complementary nature of these systems, targeting the cellulose and hemicellulose backbones (such as *Cytophaga*) or preferentially focused on the disassembly of hemicellulosic branches (such as *Asticcacaulis*), supports the existence of different ecological niches for specialist cellulolytic bacteria and their potential synergy in the decomposition of complex polysaccharides. Bacteria are also relevant members of the community involved in the degradation of microbial dead biomass in forest soil, using enzymes that are still unknown. Unlike plant biomass, composed mainly by cellulose, hemicellulose and lignin, the composition of microbial biomass appears to be much more diverse and complex. The degradation of this pool of various polysaccharides (glucans, mannans, chitin, or galactans) and glycoconjugates (glycopolymers, glycosyl phosphates or glycoproteins) composing microbial cell walls has not yet been properly addressed and deserves further attention. The importance of microbial biomass, its pool size and turnover rates in forest soils are also still difficult to quantify (Brabcová et al., 2018; Ekblad et al., 2013), and understanding the fate of dead microbial biomass in soils and its importance in C cycling deserves future attention as well. The results of this study indicating specific roles of microbial taxa in C cycling improves our potential to develop models of C cycling in terrestrial ecosystems and to improve the predictions about the fate of C in soils.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported by the Czech Science Foundation (18-26221Y) and by the project "BIOCEV - Biotechnology and Biomedicine Centre of the Academy of Sciences and Charles University CZ.1.05/1.1.00/02.0109" provided by the Ministry of Education, Youth and Sports of CR and ERDF. We acknowledge Petr Kohout for his help with the statistical analyses. The authors state that there is no conflict of interest regarding the present manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.soilbio.2020.107875>.

References

- Abe, K., Nakajima, M., Yamashita, T., Matsunaga, H., Kamisuki, S., Nihira, T., Takahashi, Y., Sugimoto, N., Miyana, A., Nakai, H., Arakawa, T., Fushinobu, S., Taguchi, H., 2017. Biochemical and structural analyses of a bacterial endo-beta-1,2-glucanase reveal a new glycoside hydrolase family. *Journal of Biological Chemistry* 292, 7487–7506.
- Alneberg, J., Bjarnason, B.S., de Bruijn, I., Schirmer, M., Quick, J., Ijaz, U.Z., Lahti, L., Loman, N.J., Andersson, A.F., Quince, C., 2014. Binning metagenomic contigs by coverage and composition. *Nature Methods* 11, 1144–1146.
- Baldrian, P., Merhautová, V., Cajthaml, T., Petráňková, M., Šnajdr, J., 2010. Small-scale distribution of extracellular enzymes, fungal, and bacterial biomass in *Quercus petraea* forest topsoil. *Biology and Fertility of Soils* 46, 717–726.
- Baldrian, P., Šnajdr, J., Merhautová, V., Dobiášová, P., Cajthaml, T., Valášková, V., 2013a. Responses of the extracellular enzyme activities in hardwood forest to soil temperature and seasonality and the potential effects of climate change. *Soil Biology and Biochemistry* 56, 60–68.
- Baldrian, P., Větrovský, T., Cajthaml, T., Dobiášová, P., Petráňková, M., Šnajdr, J., Eichlerová, I., 2013. Estimation of fungal biomass in forest litter and soil. *Fungal Ecol.* 6, 1–11.
- Bárta, J., Tahovská, K., Santrůčková, H., Oulehle, F., 2017. Microbial communities with distinct denitrification potential in spruce and beech soils differing in nitrate leaching. *Scientific Reports* 7, 9738.
- Bazaka, K., Crawford, R.J., Nazarenko, E.L., Ivanova, E.P., 2011. Bacterial extracellular polysaccharides. *Advances in Experimental Medicine & Biology* 715, 213–226.
- Berlemont, R., Martiny, A.C., 2015. Genomic potential for polysaccharides deconstruction in bacteria. *Applied and Environmental Microbiology* 81, 1513–1519.
- Bhatnagar, J.M., Peay, K.G., Treseder, K.K., 2018. Litter chemistry influences decomposition through activity of specific microbial functional guilds. *Ecological Monographs* 88, 429–444.
- Boddy, L., Hiscox, J., 2016. Fungal ecology: principles and mechanisms of colonization and competition by saprotrophic fungi. *Microbiology Spectrum* 4, 0019.
- Bomble, Y.J., Lin, C.Y., Amore, A., Wei, H., Holwerda, E.K., Ciesielski, P.N., Donohoe, B.S., Decker, S.R., Lynd, L.R., Himmel, M.E., 2017. Lignocellulose deconstruction in the biosphere. *Current Opinion in Chemical Biology* 41, 61–70.
- Brabcová, V., Nováková, M., Davidová, A., Baldrian, P., 2016. Dead fungal mycelium in forest soil represents a decomposition hotspot and a habitat for a specific microbial community. *New Phytologist* 210, 1369–1381.
- Brabcová, V., Štursová, M., Baldrian, P., 2018. Nutrient content affects the turnover of fungal biomass in forest topsoil and the composition of associated microbial communities. *Soil Biology and Biochemistry* 118, 187–198.
- Brown, S., Santa Maria Jr., J.P., Walker, S., 2013. Wall teichoic acids of gram-positive bacteria. *Annual Review of Microbiology* 67, 313–336.
- Cavaliere, M., Feng, S., Soyer, O.S., Jiménez, J.I., 2017. Cooperation in microbial communities and their biotechnological applications. *Environmental Microbiology* 19, 2949–2963.
- Cragg, S.M., Beckham, G.T., Bruce, N.C., Bugg, T.D., Distel, D.L., Dupree, P., Etxabe, A. G., Goodell, B.S., Jellison, J., McGeehan, J.E., McQueen-Mason, S.J., Schnorr, K., Walton, P.H., Watts, J.E., Zimmer, M., 2015. Lignocellulose degradation mechanisms across the tree of life. *Current Opinion in Chemical Biology* 29, 108–119.
- de Boer, W., Folman, L.B., Summerbell, R.C., Boddy, L., 2005. Living in a fungal world: impact of fungi on soil bacterial niche development. *FEMS Microbiology Reviews* 29, 795–811.
- Donohoe, B.S., Resch, M.G., 2015. Mechanisms employed by cellulase systems to gain access through the complex architecture of lignocellulosic substrates. *Current Opinion in Chemical Biology* 29, 100–107.
- Egan, A.J., Cleverley, R.M., Peters, K., Lewis, R.J., Vollmer, W., 2017. Regulation of bacterial cell wall growth. *FEBS Journal* 284, 851–867.
- Eichorst, S.A., Kuske, C.R., 2012. Identification of cellulose-responsive bacterial and fungal communities in geographically and edaphically different soils by using stable isotope probing. *Applied and Environmental Microbiology* 78, 2316–2327.
- Ekblad, A., Wallander, H., Godbold, D.L., Cruz, C., Johnson, D., Baldrian, P., Björk, R.G., Epron, D., Kieliszewska-Rokicka, B., Kjoller, R., Kraigher, H., Matzner, E., Neumann, J., Plassard, C., 2013. The production and turnover of extramatrical mycelium of ectomycorrhizal fungi in forest soils: role in carbon cycling. *Plant and Soil* 366, 1–27.
- Eren, A.M., Esen, O.C., Quince, C., Vineis, J.H., Morrison, H.G., Sogin, M.L., Delmont, T. O., 2015. Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* 3, e1319.
- Fernandez, C.W., Kennedy, P.G., 2016. Revisiting the 'Gadgil effect': do interguild fungal interactions control carbon cycling in forest soils? *New Phytologist* 209, 1382–1394.
- Fernandez, C.W., Koide, R.T., 2014. Initial melanin and nitrogen concentrations control the decomposition of ectomycorrhizal fungal litter. *Soil Biology and Biochemistry* 77, 150–157.
- Fernandez, C.W., Langley, J.A., Chapman, S., McCormack, M.L., Koide, R.T., 2016. The decomposition of ectomycorrhizal fungal necromass. *Soil Biology and Biochemistry* 93, 38–49.
- Fesel, P.H., Zuccaro, A., 2016. beta-glucan: crucial component of the fungal cell wall and elusive MAMP in plants. *Fungal Genetics and Biology* 90, 53–60.
- Fierer, N., Lauber, C.L., Ramirez, K.S., Zaneveld, J., Bradford, M.A., Knight, R., 2012. Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients. *The ISME Journal* 6, 1007–1017.
- Free, S.J., 2013. Fungal cell wall organization and biosynthesis. *Advances in Genetics* 81, 33–82.
- Fu, L., Niu, B., Zhu, Z., Wu, S., Li, W., 2012. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* 28, 3150–3152.
- Gharechahi, J., Salekdeh, G.H., 2018. A metagenomic analysis of the camel rumen's microbiome identifies the major microbes responsible for lignocellulose degradation and fermentation. *Biotechnology for Biofuels* 11, 216.
- Graham, E.B., Knelman, J.E., Schindlbacher, A., Siciliano, S., Breulmann, M., Yannarell, A., Beman, J.M., Abell, G., Philippot, L., Prosser, J., Foulquier, A., Yuste, J.C., Glanville, H.C., Jones, D.L., Angel, R., Salminen, J., Newton, R.J., Burgmann, H., Ingram, L.J., Hamer, U., Siljanen, H.M., Peltoniemi, K., Potthast, K., Baneras, L., Hartmann, M., Banerjee, S., Yu, R.Q., Nogaró, G., Richter, A., Koranda, M., Castle, S.C., Goberna, M., Song, B., Chatterjee, A., Nunes, O.C., Lopes, A.R., Cao, Y., Kaisermann, A., Hallin, S., Strickland, M.S., Garcia-Pausas, J., Barba, J., Kang, H., Isobe, K., Papaspyrou, S., Pastorelli, R., Lagomarsino, A., Lindstrom, E.S., Basiliko, N., Nemergut, D.R., 2016. Microbes as engines of ecosystem function: when does community structure enhance predictions of ecosystem processes? *Frontiers in Microbiology* 7, 214.
- Gunina, A., Dippold, M., Glaser, B., Kuz'yakov, Y., 2017. Turnover of microbial groups and cell components in soil: 13C analysis of cellular biomarkers. *Biogeosciences* 14, 271–283.
- Hamed, J., Poorinmohammad, N., 2017. The cellular structure of Actinobacteria. In: Wink, J., Mohamadippanah, F., Hamed, J. (Eds.), *Biology and Biotechnology of Actinobacteria*. Springer International Publishing, Berlin, pp. 5–28.
- Hesse, C.N., Mueller, R.C., Vuyisich, M., Gallegos-Graves, L.V., Gleason, C.D., Zak, D.R., Kuske, C.R., 2015. Forest floor community metatranscriptomes identify fungal and bacterial responses to N deposition in two maple forests. *Frontiers in Microbiology* 6, 337.
- Kang, D.D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., Wang, Z., 2019. MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* 7, e7359.

- Koh, H.W., Hur, M., Kang, M.S., Ku, Y.B., Ghai, R., Park, S.J., 2017. Physiological and genomic insights into the lifestyle of arsenite-oxidizing *Herminiimonas arsenitoxydans*. *Scientific Reports* 7, 15007.
- Kramer, S., Dibbern, D., Moll, J., Huenninghaus, M., Koller, R., Krueger, D., Marhan, S., Ulrich, T., Wubet, T., Bonkowski, M., Buscot, F., Lueders, T., Kandeler, E., 2016. Resource partitioning between bacteria, fungi, and protists in the detritusphere of an agricultural soil. *Frontiers in Microbiology* 7, 1524.
- Langmead, B., Trapnell, C., Pop, M., Salzberg, S.L., 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biology* 10, R25.
- Lasa, A.V., Masinova, T., Baldrian, P., Fernandez-Lopez, M., 2019. Bacteria from the endosphere and rhizosphere of *Quercus* spp. use mainly cell wall-associated enzymes to decompose organic matter. *Plos one* 14, e0214422.
- Levasseur, A., Drula, E., Lombard, V., Coutinho, P.M., Henrissat, B., 2013. Expansion of the enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes. *Biotechnology for Biofuels* 6, 41.
- Li, W., Godzik, A., 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* 22, 1658–1659.
- Lladó, S., Větrovský, T., Baldrian, P., 2019. Tracking of the activity of individual bacteria in temperate forest soils shows guild-specific responses to seasonality. *Soil Biology and Biochemistry* 135, 275–282.
- Lloyd, K.G., Steen, A.D., Ladau, J., Yin, J., Crosby, L., 2018. Phylogenetically novel uncultured microbial cells dominate earth microbiomes. *mSystems* 3, e00055–18.
- Lombard, V., Golconda Ramulu, H., Drula, E., Coutinho, P.M., Henrissat, B., 2014. The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Research* 42, D490–D495.
- López-Mondéjar, R., Voříšková, J., Větrovský, T., Baldrian, P., 2015. The bacterial community inhabiting temperate deciduous forests is vertically stratified and undergoes seasonal dynamics. *Soil Biology and Biochemistry* 87, 43–50.
- López-Mondéjar, R., Zühlke, D., Becher, D., Riedel, K., Baldrian, P., 2016. Cellulose and hemicellulose decomposition by forest soil bacteria proceeds by the action of structurally variable enzymatic systems. *Scientific Reports* 6, 25279.
- López-Mondéjar, R., Algora, C., Baldrian, P., 2019. Lignocellulolytic systems of soil bacteria: a vast and diverse toolbox for biotechnological conversion processes. *Biotechnology Advances* 37, 107374.
- López-Mondéjar, R., Brabcova, V., Stursova, M., Davidova, A., Jansa, J., Cajthaml, T., Baldrian, P., 2018. Decomposer food web in a deciduous forest shows high share of generalist microorganisms and importance of microbial biomass recycling. *ISME Journal* 12, 1768–1778.
- López-Mondéjar, R., Zühlke, D., Vetrovsky, T., Becher, D., Riedel, K., Baldrian, P., 2016. Decoding the complete arsenal for cellulose and hemicellulose deconstruction in the highly efficient cellulose decomposer *Paenibacillus* O199. *Biotechnology for Biofuels* 9, 104.
- Meyer, F., Paarmann, D., D'Souza, M., Olson, R., Glass, E.M., Kubal, M., Paczian, T., Rodriguez, A., Stevens, R., Wilke, A., Wilkening, J., Edwards, R.A., 2008. The metagenomics RAST server - a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* 9, 386.
- Mishra, A., Jha, B., 2013. Microbial exopolysaccharides. In: Rosenberg, E., DeLong, E.F., Stackebrandt, E., Lory, S., Thompson, F. (Eds.), *The Prokaryotes—Applied Bacteriology and Biotechnology*. Springer-Verlag, Berlin, Germany, pp. 179–192.
- Nguyen, S.T.C., Freund, H.L., Kasanjian, J., Berlemont, R., 2018. Function, distribution, and annotation of characterized cellulases, xylanases, and chitinases from CAZY. *Applied Microbiology and Biotechnology* 102, 1629–1637.
- Pan, Y., Birdsey, R.A., Fang, J., Houghton, R., Kauppi, P.E., Kurz, W.A., Phillips, O.L., Shvidenko, A., Lewis, S.L., Canadell, J.G., Ciais, P., Jackson, R.B., Pacala, S.W., McGuire, A.D., Piao, S., Rautiainen, A., Sitch, S., Hayes, D., 2011. A large and persistent carbon sink in the world's forests. *Science* 333, 988–993.
- Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W., 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Research* 25, 1043–1055.
- Parks, D.H., Rinke, C., Chuvochina, M., Chaumeil, P.A., Woodcroft, B.J., Evans, P.N., Hugenholtz, P., Tyson, G.W., 2017. Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. *Nature Microbiology* 2, 1533–1542.
- Parks, D.H., Chuvochina, M., Waite, D.W., Rinke, C., Skarshewski, A., Chaumeil, P.A., Hugenholtz, P., 2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nature Biotechnology* 36, 996–1004.
- Peng, Y., Leung, H.C., Yiu, S.M., Chin, F.Y., 2012. IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 28, 1420–1428.
- Pold, G., Billings, A.F., Blanchard, J.L., Burkhardt, D.B., Frey, S.D., Melillo, J.M., Schnabel, J., van Diepen, L.T., DeAngelis, K.M., 2016. Long-Term warming alters carbohydrate degradation potential in temperate forest soils. *Applied and Environmental Microbiology* 82, 6518–6530.
- RCORETeam, 2019. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.** <https://www.R-project.org/>.
- Rodríguez, R.L., Gunturu, S., Harvey, W.T., Rossello-Mora, R., Tiedje, J.M., Cole, J.R., Konstantinidis, K.T., 2018. The Microbial Genomes Atlas (MiGA) webservice: taxonomic and gene diversity analysis of Archaea and Bacteria at the whole genome level. *Nucleic Acids Research* 46, W282–W288.
- Rousk, J., Frey, S.D., 2015. Revisiting the hypothesis that fungal-to-bacterial dominance characterises turnover of soil organic matter and nutrients. *Ecological Monographs* 85, 457–472.
- Sahin, N., Gonzalez, J.M., Iizuka, T., Hill, J.E., 2010. Characterization of two aerobic ultramicrobacteria isolated from urban soil and a description of *Oxalicibacterium soluribus* sp. nov. *FEMS Microbiology Letters* 307, 25–29.
- Schaffer, C., Messner, P., 2005. The structure of secondary cell wall polymers: how Gram-positive bacteria stick their cell walls together. *Microbiology* 151, 643–651.
- Scheffers, D.J., Pinho, M.G., 2005. Bacterial cell wall synthesis: new insights from localization studies. *Microbiology and Molecular Biology Reviews* 69, 585–607.
- Schimel, J.P., Schaeffer, S.M., 2012. Microbial control over carbon cycling in soil. *Frontiers in Microbiology* 3, 348.
- Sieber, C.M.K., Probst, A.J., Sharrar, A., Thomas, B.C., Hess, M., Tringe, S.G., Banfield, J.F., 2018. Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nature Microbiology* 3, 836–843.
- Silhavy, T.J., Kahne, D., Walker, S., 2010. The bacterial cell envelope. *Cold Spring Harbor Perspectives in Biology* 2, a000414.
- Šnajdr, J., Valášková, V., Merhautová, V., Herinková, J., Cajthaml, T., Baldrian, P., 2008. Spatial variability of enzyme activities and microbial biomass in the upper layers of *Quercus petraea* forest soil. *Soil Biology and Biochemistry* 40, 2068–2075.
- Sommer, D.D., Delcher, A.L., Salzberg, S.L., Pop, M., 2007. Minimus: a fast, lightweight genome assembler. *BMC Bioinformatics* 8, 64.
- Štursová, M., Žifčáková, L., Leigh, M.B., Burgess, R., Baldrian, P., 2012. Cellulose utilization in forest litter and soil: identification of bacterial and fungal decomposers. *FEMS Microbiology Ecology* 80, 735–746.
- Sukdeo, N., Teen, E., Rutherford, P.M., Massicotte, H.B., Egger, K.N., 2019. Bacterial and fungal saprotrophs are strongly stimulated weeks to months after forest soil profile reconstruction. *Pedobiologia* 73, 29–41.
- Tláskal, V., Voříšková, J., Baldrian, P., 2016. Bacterial succession on decomposing leaf litter exhibits a specific occurrence pattern of cellulolytic taxa and potential decomposers of fungal mycelia. *FEMS Microbiology Ecology* 92, fiw177.
- Trivedi, P., Anderson, I.C., Singh, B.K., 2013. Microbial modulators of soil carbon storage: integrating genomic and metabolic knowledge for global prediction. *Trends in Microbiology* 21, 641–651.
- Vaaje-Kolstad, G., Forsberg, Z., Loose, J.S., Bissaro, B., Eijsink, V.G., 2017. Structural diversity of lytic polysaccharide monoxygenases. *Current Opinion in Structural Biology* 44, 67–76.
- Verastegui, Y., Cheng, J., Engel, K., Kolczynski, D., Mortimer, S., Lavigne, J., Montalibet, J., Romantsov, T., Hall, M., McConkey, B.J., Rose, D.R., Tomashek, J.J., Scott, B.R., Charles, T.C., Neufeld, J.D., 2014. Multisubstrate isotope labeling and metagenomic analysis of active soil bacterial communities. *MBio* 5, e01157-01114.
- Voříšková, J., Brabcová, V., Cajthaml, T., Baldrian, P., 2014. Seasonal dynamics of fungal communities in a temperate oak forest soil. *New Phytologist* 201, 269–278.
- Wallenstein, M.D., McNulty, S., Fernandez, I.J., Boggs, J., Schlesinger, W.H., 2006. Nitrogen fertilization decreases forest soil fungal and bacterial biomass in three long-term experiments. *Forest Ecology and Management* 222, 459–468.
- Wang, X., Sharp, C.E., Jones, G.M., Grasby, S.E., Brady, A.L., Dunfield, P.F., 2015. Stable-isotope probing identifies uncultured Planctomycetes as primary degraders of a complex heteropolysaccharide in soil. *Applied and Environmental Microbiology* 81, 4607–4615.
- Wang, X., Han, Q., Chen, G., Zhang, W., Liu, W., 2017. A putative type II secretion system is involved in cellulose utilization in *Cytophaga hutchinsonii*. *Frontiers in Microbiology* 8, 1482.
- Weidenmaier, C., Peschel, A., 2008. Teichoic acids and related cell-wall glycopolymers in Gram-positive physiology and host interactions. *Nature Reviews Microbiology* 6, 276–287.
- West, P.T., Probst, A.J., Grigoriev, I.V., Thomas, B.C., Banfield, J.F., 2018. Genome-reconstruction for eukaryotes from complex natural microbial communities. *Genome Research* 28, 569–580.
- Wilhelm, R.C., Singh, R., Ertis, L.D., Mohn, W.W., 2019. Bacterial contributions to delignification and lignocellulose degradation in forest soils with metagenomic and quantitative stable isotope probing. *ISME Journal* 13, 413–429.
- Woodward, S., Boddy, L., 2008. Interactions between saprotrophic fungi. *British Mycological Society Symposium Series* 28, 125–141.
- Wu, Y.W., Simmons, B.A., Singer, S.W., 2015. MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* 32, 605–607.
- Yin, Y., Mao, X., Yang, J., Chen, X., Mao, F., Xu, Y., 2012. dbCAN: a web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Research* 40, W445–W451.
- Zerbin, D.R., Birney, E., 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Research* 18, 821–829.
- Zhang, H., Yohe, T., Huang, L., Entwistle, S., Wu, P., Yang, Z., Busk, P.K., Xu, Y., Yin, Y., 2018. dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Research* 46, W95–W101.
- Zhu, Y., Han, L., Hefferon, K.L., Silvaggi, N.R., Wilson, D.B., McBride, M.J., 2016. Periplasmic *Cytophaga hutchinsonii* endoglucanases are required for use of crystalline cellulose as the sole source of carbon and energy. *Applied and Environmental Microbiology* 82, 4835–4845.
- Žifčáková, L., Větrovský, T., Howe, A., Baldrian, P., 2016. Microbial activity in forest soil reflects the changes in ecosystem properties between summer and winter. *Environmental Microbiology* 18, 288–301.
- Žifčáková, L., Větrovský, T., Lombard, V., Henrissat, B., Howe, A., Baldrian, P., 2017. Feed in summer, rest in winter: microbial carbon utilization in forest topsoil. *Microbiome* 5, 122.